

**MINISTRY OF PUBLIC HEALTH OF UKRAINE
NATIONAL UNIVERSITY OF PHARMACY**

Educational degree master Training area 22 – PUBLIC HEALTH
(level of educational degree) (code number and title of training area)
 Speciality 226 – PHARMACY, INDUSTRIAL PHARMACY Educational program PHARMACY (ФМ(5,0)АНГЛ)
(code number and title of speciality) (title of educational program)
 Semester spring semester, 2019/2020 academic year Subject ANALYTICAL CHEMISTRY
(title of academic subject)

THEMATIC CONTROL №4

QUESTION CARD (EXAMPLE)

- Detection of glucose and maltose has been carried out by the method of thin-layer chromatography. The next distances have been passed by glucose and maltose – 3.8 cm and 2.3 cm respectively. The solvent has passed the distance of 10.0 cm for the same time. Calculate the values of R_f for each substance to be determined.
- Determination of the substance by the method of spectrophotometry. At $\lambda = 410$ nm the molar absorption coefficient is 8000 L/mole·cm, the specific absorption coefficient is 147.75 1/%·cm. Calculate the molar mass of the substance to be determined.
- Determination of glucose in the solution by the method of polarimetry. The rotation angle of polarization plane for the solution to be analysed is $+11.80^\circ$, the layer thickness is 1 dm and the value of the specific rotation is $+53.1^\circ$. Calculate the concentration of glucose (g/100 mL) in the solution to be analysed.
- Carry out potentiometric determination of $\text{Na}_2\text{C}_2\text{O}_4$ ($M(\text{Na}_2\text{C}_2\text{O}_4) = 134.000$ g/mole) by the method of permanganatometry (the method of separate samples). Write the equation of reactions. Choose the pair of electrodes for the determination. Calculate the stoichiometrical ratio s , the factor of equivalence f for the substance to be determined and its molar mass of equivalent E . Calculate the percentage of the substance to be determined in three ways – according to the molar mass of equivalent, according to the molar mass and according to the titre of the titrant by the substances to be determined ($c(1/5\text{KMnO}_4) = 0.1015$ mole/dm³, $V(\text{KMnO}_4) = 25.18$ cm³, $m(\text{Na}_2\text{C}_2\text{O}_4) = 0.2458$ g).
- Answer the tests.

POINTS DISTRIBUTION

question 1	2 points
question 2	2 points
question 3	2 points
question 4	2 points
question 5	1 points
in all	9 points

Estimation scale: national and ECTS

Points in all	ECTS mark	Mark by national scale
8.1 – 9.0	A	5
7.3 – 8.0	B	4
6.9 – 7.3	C	
6.0 – 6.8	D	3
5.0 – 5.9	E	
0 – 4.9	F	2

It has been approved at the meeting of the Analytical Chemistry Department.
The minutes №1 from 29. 08. 2019 year.

Head of the Analytical Chemistry Department, prof. _____

(sign)

I. S. Grytsenko

Examiner, as. prof. _____

(sign)

L. Yu. Klimenko

performed by as. prof. Klimenko L. Yu., as. prof. Mykytenko O. Ye., as. prof. Kostina T. A.

THEMATIC CONTROL №4

1. Detection of glucose and maltose has been carried out by the method of thin-layer chromatography. The next distances have been passed by glucose and maltose – 3.8 cm and 2.3 cm respectively. The solvent has passed the distance of 10.0 cm for the same time. Calculate the values of R_f for each substance to be determined.

Given:

$S(\text{glucose}) = 3.8 \text{ cm}$ $S(\text{maltose}) = 2.3 \text{ cm}$ $S(\text{solvent}) = 10.0 \text{ cm}$	$R_f(\text{glucose}) = \frac{S(\text{glucose})}{S(\text{solvent})} = \frac{3.8}{10.0} = 0.38$
$R_f(\text{glucose}) - ?$ $R_f(\text{maltose}) - ?$	$R_f(\text{maltose}) = \frac{S(\text{maltose})}{S(\text{solvent})} = \frac{2.3}{10.0} = 0.23$

2. Determination of the substance by the method of spectrophotometry. At $\lambda = 410 \text{ nm}$ the molar absorption coefficient is $8000 \text{ L/mole}\cdot\text{cm}$, the specific absorption coefficient is $147.75 \text{ 1/\%}\cdot\text{cm}$. Calculate the molar mass of the substance to be determined.

Given:


$\epsilon = 8000 \text{ L/mole}\cdot\text{cm}$ $A_{1\text{cm}}^{1\%} = 147.75 \text{ 1/\%}\cdot\text{cm}$	$M = \frac{\epsilon}{A_{1\text{cm}}^{1\%}} \cdot 10 = \frac{8000}{147.75} \cdot 10 = 541.46 \text{ g/mole}$
$M - ?$	

3. Determination of glucose in the solution by the method of polarimetry. The rotation angle of polarization plane for the solution to be analysed is $+11.80^\circ$, the layer thickness is 1 dm and the value of the specific rotation is $+53.1^\circ$. Calculate the concentration of glucose ($\text{g}/100 \text{ mL}$) in the solution to be analysed.

Given:

$\alpha = +11.80^\circ$ $[\alpha]_D^{20} = +53.1^\circ$	$C = \frac{\alpha \cdot 100}{[\alpha]_D^{20} \cdot l} = \frac{11.80 \cdot 100}{53.1 \cdot 1} = 22.2 \text{ g}/100 \text{ mL}$
$C - ?$	

4. Carry out potentiometric determination of $\text{Na}_2\text{C}_2\text{O}_4$ ($M(\text{Na}_2\text{C}_2\text{O}_4) = 134.000 \text{ g/mole}$) by the method of permanganatometry (the method of separate samples). Write the equation of reactions. Choose the pair of electrodes for the determination. Calculate the stoichiometrical ratio s , the factor of equivalence f for the substance to be determined and its molar mass of equivalent E . Calculate the percentage of the substance to be determined in three ways – according to the molar mass of equivalent, according to the molar mass and according to the titre of the titrant by the substances to be determined ($c(1/5\text{KMnO}_4) = 0.1015 \text{ mole/dm}^3$, $V(\text{KMnO}_4) = 25.18 \text{ cm}^3$, $m(\text{Na}_2\text{C}_2\text{O}_4) = 0.2458 \text{ g}$).



$$s = 5/2; f(\text{Na}_2\text{C}_2\text{O}_4) = 1/2; f(\text{KMnO}_4) = 1/5;$$

$$E(\text{Na}_2\text{C}_2\text{O}_4) = M(\text{Na}_2\text{C}_2\text{O}_4) \cdot f(\text{Na}_2\text{C}_2\text{O}_4) = 134.000 \cdot 1/2 = 67.000 \text{ g/mole}$$

Given:

$$c(1/5\text{KMnO}_4) = 0.1015 \text{ mole/dm}^3$$

$$V(\text{KMnO}_4) = 25.18 \text{ cm}^3$$

$$m(\text{Na}_2\text{C}_2\text{O}_4) = 0.2458 \text{ g}$$

$$E(\text{Na}_2\text{C}_2\text{O}_4) = 67.000 \text{ g/mole}$$

$$M(\text{Na}_2\text{C}_2\text{O}_4) = 134.000 \text{ g/mole}$$

$$\omega(\text{Na}_2\text{C}_2\text{O}_4) - ?$$

calculation of $\omega(\text{Na}_2\text{C}_2\text{O}_4)$ according to $E(\text{Na}_2\text{C}_2\text{O}_4)$

$$\begin{aligned} \omega(\text{Na}_2\text{C}_2\text{O}_4) &= \frac{c(1/5\text{KMnO}_4) \cdot V(\text{KMnO}_4) \cdot E(\text{Na}_2\text{C}_2\text{O}_4) \cdot 100}{1000 \cdot m(\text{Na}_2\text{C}_2\text{O}_4)} = \\ &= \frac{0.1015 \cdot 25.18 \cdot 67.000 \cdot 100}{1000 \cdot 0.2458} = 69.70\% \end{aligned}$$

$$c(\text{KMnO}_4) = c(1/5\text{KMnO}_4) \cdot f(\text{KMnO}_4) = 0.1015 \cdot 1/5 = 0.02030 \text{ mole/dm}^3$$

calculation of $\omega(\text{Na}_2\text{C}_2\text{O}_4)$ according to $M(\text{Na}_2\text{C}_2\text{O}_4)$ and s

$$\begin{aligned} \omega(\text{Na}_2\text{C}_2\text{O}_4) &= \frac{c(\text{KMnO}_4) \cdot V(\text{KMnO}_4) \cdot s \cdot M(\text{Na}_2\text{C}_2\text{O}_4) \cdot 100}{1000 \cdot m(\text{Na}_2\text{C}_2\text{O}_4)} = \\ &= \frac{0.02030 \cdot 25.18 \cdot 5/2 \cdot 134.000 \cdot 100}{1000 \cdot 0.2458} = 69.70\% \end{aligned}$$

calculation of $\omega(\text{Na}_2\text{C}_2\text{O}_4)$ according to $T(\text{KMnO}_4/\text{Na}_2\text{C}_2\text{O}_4)$

$$T(\text{KMnO}_4/\text{Na}_2\text{C}_2\text{O}_4) = \frac{c(1/5\text{KMnO}_4)_{\text{theor}} \cdot E(\text{Na}_2\text{C}_2\text{O}_4)}{1000} = \frac{0.1000 \cdot 67.000}{1000} = 0.006700 \text{ g/cm}^3$$

$$T(\text{KMnO}_4/\text{Na}_2\text{C}_2\text{O}_4) = \frac{c(\text{KMnO}_4)_{\text{theor}} \cdot s \cdot M(\text{Na}_2\text{C}_2\text{O}_4)}{1000} = \frac{0.02000 \cdot 5/2 \cdot 134.000}{1000} = 0.006700 \text{ g/cm}^3$$

$$K(\text{KMnO}_4) = \frac{c(1/5\text{KMnO}_4)_{\text{pract}}}{c(1/5\text{KMnO}_4)_{\text{theor}}} = \frac{0.1015}{0.1000} = 1.015$$

$$\begin{aligned} \omega(\text{Na}_2\text{C}_2\text{O}_4) &= \frac{K(\text{KMnO}_4) \cdot V(\text{KMnO}_4) \cdot T(\text{KMnO}_4/\text{Na}_2\text{C}_2\text{O}_4) \cdot 100}{m(\text{Na}_2\text{C}_2\text{O}_4)} = \\ &= \frac{1.015 \cdot 25.18 \cdot 0.006700 \cdot 100}{0.2458} = 69.70\% \end{aligned}$$